

**Supplementary Material for Integrin Mechanosensing relies on Pivot-clip  
Mechanism to Reinforce Cell Adhesion**

Andre R. Montes, Anahi Barroso, Wei Wang, Grace D. O'Connell, Adrian Buganza Tepole, Mohammad R.K. Mofrad

Table S1: Steered Molecular Dynamics Parameters

Parameter	Setting
Time step	2fs
Number of steps (Time) for 1nm/ns	10000000 (20ns)
Number of steps (Time) for 10nm/ns	1500000 (3ns)
Integrator	Leapfrog algorithm
Constraint Algorithm	LINEar Constraint Solver (LINCS)
Constraints	H-bonds constrained
Cutoff scheme	Verlet (Buffered neighbor searching)
Short-range neighbor list cutoff	1.4nm
Short-range electrostatic cutoff	1.4nm
Short-range van der Waals cutoff	1.4nm
Electrostatics	Fast smooth Particle-Mesh Ewald (SPME)
Interpolation order	Cubic
Grid spacing for fast Fourier Transform	0.12nm
Temperature coupling	Nosé-Hoover
Reference temperature	310K
Temperature time constant	1.0ps
Temperature coupled groups	Protein and non-protein
Pressure coupling	Off
Dispersion correction	long range dispersion corrections for energy and pressure
Velocity generation	Off
Harmonic potential	Umbrella
Force constant	50 kJ/mol-nm <sup>2</sup>
Pull direction	y-direction (vertical)
Pull rate for 1nm/ns	0.001nm/ps = 1nm/ns
Pull rate for 10nm/ns	0.010nm/ps = 10nm/ns

Table S2: Time resolved Force Distribution Analysis Parameter Settings

Parameter	Setting
Pairwise forces	Summed
Pairwise groups	Protein
Residue based calculation	Punctual Stress
Pairwise force type	Coulombic interactions only

Table S3: Catch bond parameters for whole-cell finite element model

Variable	Wildtype	R1374/9A
$K_{on}$	$0.1 \text{ s}^{-1}$	$0.02 \text{ s}^{-1}$
$K_a$	$0.4 \text{ s}^{-1}$	$0.8 \text{ s}^{-1}$
$K_b$	$4E - 7\text{s}^{-1}$	$8E - 7\text{s}^{-1}$
$F_a$		$-25\text{pN}$
$F_b$		$-15\text{pN}$




Figure S1: Root-mean-square deviation (RMSD) of wildtype and mutant during NPT simulation




Figure S2: Force and COM extension over time plotted over punctual stress at the synergy site [1373-1380] for A) 10nm/ns wildtype  $\alpha_5\beta_1$ -FN, B) 1nm/ns wildtype  $\alpha_5\beta_1$ -FN, C) 10nm/ns R1374/9A  $\alpha_5\beta_1$ -FN, and D) 1nm/ns R1374/9A  $\alpha_5\beta_1$ -FN. Positions (i), (ii), and (iii) correspond to the time at the peak force, local minimum, and final frame, respectively.